

Proof of the Morse conjecture for analytic flows on orientable surfaces

S. Aranson E. Zhuzhoma

Abstract

In 1946, M. Morse [14] proposed a conjecture that an analytic topologically transitive system is metrically transitive. We prove this Morse conjecture for flows on a closed orientable surface of negative Euler characteristic. As a consequence, the Morse conjecture is true for highly transitive analytic flows on closed non-orientable surfaces.

1 Introduction

Let M be a smooth Riemannian manifold. The Riemann structure on M induces a corresponding *Lebesgue measure* μ with a smooth density function (in particular, μ is positive on any open set of M), see exm. [16], ch. 3. Below, M is a closed manifold (compact and without boundary) and we'll assume that μ is normal.

A dynamical system on M is called *transitive*, if it has a dense orbit. A dynamical system $\mathcal{D}$ is called *metrically transitive* if, given any compact invariant set E , either E or its complement is a zero measure set with respect to μ . Note that μ is not in general an invariant measure of $\mathcal{D}$. Recall that E is said to be invariant under $\mathcal{D}$, if the whole orbit $l(x)$ through x belongs to E for every point $x \in E$.

It is known that a metrically transitive dynamical system is transitive [6]. The converse in general fails. Morse [14] conjectured that the converse is true for analytic dynamical systems $\mathcal{D}$ or $\mathcal{D}$ with some degree of analytic regularity. Here we prove this Morse conjecture for analytic flows on a closed orientable surface of negative Euler characteristic.

Let us mention three papers concerned the subject. Ding [8] proved the Morse conjecture for analytic flows on the torus T^2 (orientable closed surface

of zero Euler characteristic) and constructed a transitive C^∞ flow which is not metrically transitive on any closed n -manifold. In [9], Ding proved that a transitive C^1 flow with finitely many fixed points on a closed orientable surface is metrically transitive. This last result was proved by Marzougui [13] for flows with a countable set of fixed points. Note that an analytic flow can have an uncountable set of fixed points.

Our main result is the following theorem.

Theorem 1 *Let M^2 be a closed orientable surface of negative Euler characteristic and f^t be a transitive analytic flow on M^2 . Then f^t is metrically transitive.*

It is well-known that if a closed orientable surface M^2 admits a transitive flow, then M^2 has a non-positive Euler characteristic (see exm., [3]). Taking into account the paper [8], as a consequence, we get the following result.

Theorem 2 *Let M^2 be a closed orientable surface and f^t be a transitive analytic flow on M^2 . Then f^t is metrically transitive.*

One can show that there is a transitive flow on a non-orientable closed surface such that its double covering flow on the corresponding orientable surface is not a transitive flow. On the other hand, the double covering flow for a highly transitive flow (recall that a flow is highly transitive if every its one-dimensional trajectory is dense in M^2 [10]) is transitive. Moreover, if a non-orientable closed surface admits a highly transitive flow, then its Euler characteristic is less or equal to -2. Therefore, theorem 1 implies the following result.

Theorem 3 *Let M^2 be a closed non-orientable surface and f^t be a highly transitive analytic flow on M^2 . Then f^t is metrically transitive.*

Acknowledgment. The research was partially supported by RFFI grant 02-01-00098.

2 Previous results

For references, we formulate as a theorem the description of fixed points set of an analytic surface flow from Anosov's paper [2], p. 38-41.

Theorem 4 *Let $\text{Fix } f^t$ be a set of fixed points of analytic flow f^t on compact surface M^2 . Then $\text{Fix } f^t$ contains a finitely many isolated points and finitely many isolated simple closed curves (the union of all this isolated points and curves is denoted by Iso). The remainder $\text{Fix } f^t - \text{Iso}$ contains a finitely many points N that divide $\text{Fix } f^t - \text{Iso}$ into pairwise disjoint analytic arcs with endpoints in N . Moreover, given any arc $A \subset \text{Fix } f^t - \text{Iso}$ and given any point $a \in A - N$, there is a neighborhood $U(a)$ of the point a such that*

- *The restriction of f^t on $U(a) - A$ is topologically equivalent to a linear fixed point free flow of the type*

$$\dot{x} = 1, \quad \dot{y} = 0.$$

- *There is an arc $\Sigma \subset U(a)$ such that $\Sigma \cap A = a$ and either $\Sigma - a$ is transversal to the flow f^t or $\Sigma - a$ belongs to a one-dimensional trajectory of f^t .*

Corollary 1 *The set $\text{Fix } f^t$ of fixed points of analytic flow f^t on M^2 has a zero Lebesgue measure.*

We recall that a semitrajectory or trajectory is called a *nontrivially $\omega(\alpha)$ -recurrent* if it is contained in its $\omega(\alpha)$ -limit set, and it is neither a fixed point nor a periodic trajectory. A trajectory is *nontrivially recurrent* if it is both ω - and α -recurrent. A closure of a nontrivially $\omega(\alpha)$ -recurrent semitrajectory is called a *quasiminimal set*. Cherry [7] proved that a quasiminimal set contains a continuum nontrivially recurrent trajectories each of them is dense in the quasiminimal set (this result is true in a paracompact space). It is not true in general that every nontrivially recurrent trajectory is dense in the quasiminimal set. However, Maier [12] proved that this result is correct for flows on compact surfaces. Moreover, he has got a criterion of nontrivial recurrentness and proved a mutual limiting of nontrivially recurrent semitrajectories. Formally, Maier assumed that a surface flow has finitely many fixed points, but as mentioned in [4], Maier's proofs are valid for flows with any set of fixed points (the finiteness of fixed points set Maier used for other results, see the sketch of proofs of Maier's theorems in [4], [15]). We represent these Maier's results as the following theorem.

Theorem 5 *Let f^t be a flow (possibly, topological) on a closed orientable surface M^2 . Then*

1. If one-dimensional semitrajectory l belongs to the limit set of some semitrajectory and the limit set of l contains at least one point that is not a fixed point, then l is a nontrivially recurrent semitrajectory.
2. If a nontrivially recurrent semitrajectory l_1 belongs to the limit set of a nontrivially recurrent semitrajectory l_2 , then l_2 belongs to the limit set of l_1 .

Following [1], let us give the definition of ω -separatrix (the definition of α -separatrix is similar). Let $l^+(m_0) = l^+$ be a positive one-dimensional semitrajectory with the ω -limit set $\omega(l^+)$ being a unique fixed point, say s . Let Σ be a transversal segment through m_0 . Suppose that there are a neighborhood¹ $U(s)$ and a sequence of points m_k such that

1. $m_k \rightarrow m_0$ as $k \rightarrow \infty$, where $m_k \in \Sigma - m_0$.
2. $m_0 \notin U(s)$, $m_k \notin U(s)$.
3. Starting with m_k , the positive semitrajectory $l_k^+(m_k)$ enters $U(s)$ and after leaves $U(s)$.

Such l^+ is called an ω -separatrix with respect to $U(s)$. If l^+ is the ω -separatrix with respect to any sufficiently small neighborhood of s , then l^+ is called a (simply) ω -separatrix.

A *separatrix connection* is both an ω - and α -separatrix. A *separatrix loop* l is a particular case of separatrix connection, when $\omega(l) = \alpha(l)$.

Below, one considers analytic flows on a closed orientable surface M^2 , unless otherwise stated.

Lemma 1 *Let f^t be a transitive analytic flow on M^2 . Then*

1. Every separatrix with respect to some neighborhood is a (simply) separatrix.
2. If l is a positive (negative) one-dimensional semitrajectory such that the $\omega(\alpha)$ -limit set of l is a unique fixed point, then l is an $\omega(\alpha)$ -separatrix.
3. f^t has a finitely many separatrix connections.

¹Without loss of generality, one can assume that the closure of $U(s)$ is homeomorphic to a closed disk.

Proof. Due to the transitivity of f^t , every separatrix with respect to some neighborhood is a (simply) separatrix. The second assertion follows immediately from the transitivity and definition of a separatrix. Assume that the third assertion is not correct. Hence, due to M^2 has a finite genus, there is a simply connected domain bounded by separatrix connections and arcs (possibly, trivial) that belongs to $Fix f^t$. This contradicts to the transitivity of f^t and the Poincare-Bendixon theorem. $\square$

Following [11], let us give the definition of a Σ -arc. Let Σ be a transversal segment intersected by a trajectory l at the points $a, b \in l \cap \Sigma$. Denote by $\widehat{ab}$ the arc of l between a, b . Such an arc is called a Σ -arc if it has no intersections with Σ except the points a, b . The segment $[a, b] \subset \Sigma$ is called a Σ -base of Σ -arc $\widehat{ab}$. Clearly, $[a, b] \cup \widehat{ab}$ is a closed simple curve called a Σ -loop.

Suppose that the intersection $l \cap \Sigma$ consists of a countable set of points. Then there is a countable set of Σ -arcs generated by l . Consider some sequence $\widehat{a_n b_n}$ of Σ -arcs.

Lemma 2 *If Σ -base $[a_n b_n]$ are pairwise disjoint, then, beginning with some subscript, consecutive Σ -arcs bound an annulus on M^2 .*

Proof follows from lemma 1 [12], which is actually a consequence of the fact that the surface M^2 has a finite genus (see also lemma 2.8 [3]). $\square$

3 Proof of main theorem

Before the proving of main theorem 1, we consider a series of lemmas.

Let Σ be a segment endowed with a one-to-one surjective parametrization $[0, 1] \rightarrow \Sigma$ that defines a natural order relation between points of Σ : a point $a \in \Sigma$ is less than a point $b \in \Sigma$, if the parameter of a is less than the parameter of b . This ordering induces the order relation between any disjoint intervals of Σ (I is less than J , where $I, J \subset \Sigma$, $I \cap J = \emptyset$, if any point of I is less than any point of J). A countable family of pairwise disjoint intervals is a *monotone sequence*, if every current interval is less than the consecutive interval.

Any trajectory has the natural time parametrization. Therefore one can define similarly the order relation between disjoint arcs and a monotone sequence of pairwise disjoint arcs (in particular, Σ -arcs) of the trajectory.

Assume that the interior of Σ is transversal to a flow and a trajectory l intersects Σ at infinitely many points. Thus, there are infinitely many Σ -arcs.

Suppose that one of the endpoints of Σ , say c , is an accumulation point for the set $\Sigma \cap l$. One can assume that c corresponds to the parameter 1 and another endpoint b of Σ corresponds to 0. In such a notation, the following lemma takes place.

Lemma 3 *At least one of the following cases takes place: 1) there is a sequence of pairwise disjoint Σ -loops whose Σ -bases form a monotone sequence converging to c ; 2) the intersection $\Sigma \cap l$ has an accumulation point on $\Sigma - c$.*

Proof. Take some Σ -arc $\widehat{a_1 b_1}$ and put $c_1 = \max\{a_1, b_1\}$. Since c is an accumulation point for the set $\Sigma \cap l$, there are Σ -arc that is greater than $\widehat{a_1 b_1}$ and begins on (c_1, c) . If any of such Σ -arc ends on (b, c_1) , then the intersection $\Sigma \cap l$ has an accumulation point on $\Sigma - c$ (more exactly, on $\Sigma - (c, c_1)$). Suppose that there is a Σ -arc $\widehat{a_2 b_2}$ that is greater than $\widehat{a_1 b_1}$ and begins on (c_1, c) , and ends on (c_1, c) . Put $c_2 = \max\{a_2, b_2\}$. Again, if any Σ -arc that is greater than $\widehat{a_2 b_2}$ and begins on (c_2, c) ends on (b, c_2) , then the intersection $\Sigma \cap l$ has an accumulation point on $\Sigma - c$. Continuing in such a way, we either get a sequence of Σ -loops satisfying case 1), or the process will break and we get case 2). $\square$

Lemma 4 *Suppose that case 1) of lemma 3 holds, and let f^t is transitive. Then there is an interval $I \subset \Sigma - c$ with the endpoint c such that I belongs to a domain of Poincare map induced by f^t .*

Proof. Consider the set S of first intersections of ω -separatrices with $\Sigma - c$,

$$S = \{x \in \Sigma - c \mid l^+(x) \text{ is an } \omega - \text{separatrix and } l^+(x) \cap \Sigma = \{x\}\}.$$

First, let us prove that there is a nontrivial interval $I_0 \subset \Sigma - c$ with the endpoint c such that $I_0 \cap S = \emptyset$. Suppose the contrary. Then there is a sequence of points $x_n \in S$ such that $x_n \rightarrow c$ as $n \rightarrow \infty$. It follows from theorem 4 that the set $Fix f^t$ consists of a finitely many arcwise connected component. Hence for some number $n \in \mathbb{N}$, ω -separatrices $l^+(x_n)$, $l^+(x_{n+1})$ and the segment $[x_n, x_{n+1}] \subset \Sigma$, and the fixed points $\omega(l^+(x_n))$, $\omega(l^+(x_{n+1}))$, and an arc connecting this points and belonging $Fix f^t$ bound a simply connected domain in M^2 (so-called, a generalized Bendixon's sack). Therefore any positive semi-trajectory entering in this domain can't leave it. This contradicts a transitivity of f^t . Thus, we prove the existence of the nontrivial interval $I_0 \subset \Sigma - c$ with the endpoint c such that $I_0 \cap S = \emptyset$.

Since case 1) takes place, there are nontrivial intervals on I_0 that belong to a domain of Poincare map. According to lemma 3.7 [3], the endpoints of a maximal interval of Poincare map belong to ω -separatrices. Moreover, after the endpoints these ω -separatrices have no intersections with $\Sigma - c$. Hence, $I_0 = I$ is in a domain of Poincare map. $\square$

Lemma 5 *Suppose that case 1) of lemma 3 holds. Then f^t is not transitive.*

Proof. Let $I \subset \Sigma - c$ be the interval satisfying lemma 4. Without loss of generality, one can assume that Σ -bases of Σ -loops of l belong to I . Moreover, one can assume that the Σ -loops satisfy to lemma 2, so every consecutive Σ -loops bound an annulus on M^2 . Let us show that the union of this annuluses (denoted by K) is an open annulus.

Glue artificially a closed disk D to the first Σ -loop. Since Σ -bases are pairwise disjoint, the corresponding annuluses are pairwise disjoint as well. Therefore the union of annuluses one can represent as the union of nested increasing disks. It is known that such the union is an open disk. Removing D , we see that the union K of annuluses is an open annulus.

By construction, $I \subset K$. Due to lemma 4, any positive semitrajectory starting on I must intersect I . Moreover, the arc of such a positive semitrajectory between intersections with I belongs to K because K is a union of annuluses formed by Σ -loops. Therefore any positive semitrajectory that enter in K can't leave K . Since M^2 is not a torus, $\text{clos } K \neq M^2$. This contradicts to a transitivity of f^t . $\square$

Lemma 6 *Suppose that case 2) of lemma 3 holds and f^t is transitive. Then the semitrajectory l is dense on M^2 .*

Proof. By condition of case 2), there is a point, say $z \in \Sigma - c$, that belongs to ω -limit set $\omega(l)$ of the semitrajectory l . Since $z \in \Sigma - c$, z is not a fixed point. By the transitivity of f^t , l is in ω -limit set of some semitrajectory, which is dense on M^2 . According to theorem 5, l is a nontrivially recurrent semitrajectory and is dense on M^2 . $\square$

Lemma 7 *Let N be a closed invariant set of transitive analytic flow f^t on M^2 , and suppose that $N \neq M^2$. Then any one-dimensional non-periodic trajectory $l \subset N$ is a separatrix connection.*

Proof. Due to lemma 1, it is sufficient to prove that both the ω - and α -limit set of l is a unique fixed point. We'll consider the only ω -limit set $\omega(l)$, the proof for $\alpha(l)$ is similar.

Suppose the contrary. Obviously, $\omega(l)$ is non-empty. Then at least one of the following cases take place: a) $\omega(l)$ contains continuum fixed points; b) $\omega(l)$ contains a point that is not fixed (so-called, a regular point). In the both cases, taking into account theorem 4, there exists a transversal segment Σ (possibly, open) intersected by l infinitely many times. Moreover, one of the endpoints of Σ , say c , is in $\omega(l)$ and is an accumulation point of the intersection $l \cap \Sigma$. Note that c could be either a fixed point or regular one. We see that one of the two cases of lemma 3 holds. Due to lemma 5 and a transitivity of f^t , one holds case 2) of lemma 3. Hence, according to lemma 6, l is dense on M^2 . This contradicts to the condition $N \neq M^2$. $\square$

Proof of theorem 1. Let N be a closed invariant set of the transitive analytic flow f^t on M^2 . For $N = M^2$, there is nothing to prove. Suppose that $N \neq M^2$. Then corollary 1 and lemmas 1, 7 imply that N has a zero Lebesgue measure. Hence, f^t is metrically transitive. $\square$

References

- [1] **Andronov A. A., Leontovich E. A., Gordon I. I., Maier A.G.** *Qualitative theory of dynamical systems of the second order.* **1966**, Moscow (Russian); MR 33# 7650. *English transl.*: Halstead Press, New York-Toronto; Israel Program for Sci. Translations, Jerusalem-London, 1973, p. 524. MR 50# 2619.
- [2] **Anosov D. V.** On the behavior of trajectories, on the Euclidian and Lobachevsky plane, covering trajectories of flows on closed surfaces. I, *Izvestia Acad. Nauk SSSR, Ser. Mat.* **51**(1987), no 1, 16-43 (in Russian); *Transl. in: Math. USSR, Izv.* **30**(1988); MR 88j:58111.
- [3] **Aranson S., Belitsky G., Zhuzhoma E.** *An Introduction to Qualitative Theory of Dynamical Systems on Surfaces.* Amer. Math. Soc., Math. Monogr., Providence, **1996**.
- [4] **Aranson S., Zhuzhoma E.** Maier's theorems and geodesic laminations of surface flows. *Journ. of Dyn. and Contr. Syst.*, **2**(1996), no 4, 557-582.

- [5] **Bendixson I.** Sur les courbes définies par les equations différentielles. *Acta Math.* **24**(1901), 1-88.
- [6] **Birkhoff G. D.** *Dynamical systems*, AMS, New York, **1927**.
- [7] **Cherry T.** Topological properties of solutions of ordinary differential equations. *Amer. J. Math.* **59**(1937), 957-982.
- [8] **Ding T.** On Morse conjecture of metric transitivity. *Sci. China*, ser. A, **34**(1991), 2, 138-146.
- [9] **Ding T.** An ergodic theorem for flows on closed surfaces. *Nonlinear Anal.*, **35**(1999), 669-676.
- [10] **Gardiner C.J.** The structure of flows exhibiting nontrivial recurrence on two-dimensional manifolds. *Journ. Diff. Equat.*, 1985, **57**, 1, 138-158.
- [11] **Gutierrez C.** Structural stability for flows on the torus with a cross-cap. *Trans. AMS*, **241**(1978), 311-320.
- [12] **Maier A.** On trajectories on closed orientable surfaces, *Mat. Sb.*, **12**(1943), 71-84 (Russian), MR 5-156.
- [13] **Marzougui H.** On Morse conjecture for flows on closed surfaces. *Math. Nachr.*, **241**(2002), 121-124.
- [14] **Morse M.** George David Birkhoff and his mathematical works. *Bull. Amer. Math. Soc.*, **52**(1946), 5, 357-391.
- [15] **Nikolaev I., Zhuzhoma E.** Flows on 2-dimensional manifolds. *Lect. Notes* **1705**(1999).
- [16] **Sternberg S.** *Lectures on Differential Geometry*. Printice Hall, N.J., **1964**.

2875 Cowley Way (1015), San Diego, CA 92110, USA

E-mail: saranson@yahoo.com

Nizhny Novgorod State Technical University, Russia

E-mail: zhuzhoma@mail.ru